

D-rise 200000IU/ml injection safety card

Dosage and administration:

1 ampoule contains 200,000 IU of vitamin D3 Administered intramuscularly

Precautions :

- To avoid overdose, take into account the total doses of vitamin D when combining several treatments containing this vitamin.
- Monitor calciuria and serum calcium and stop vitamin D intake if serum calcium exceeds 105 mg / ml (2.62 mmol / l) or if calciuria exceeds 4 mg / kg / day in adults or 4-6 mg / kg / day in children.
- In case of high intake of calcium, a regular check of the calciurie is essential.

Contraindication :

- Hypersensitivity to the active substance or to any of the excipients .
- Hypercalcemia, hypercalciuria, calcium lithiasis.

Pregnancy and lactation:

There is no teratogenesis study available in animals. Clinically, a significant decline seems to exclude a malformative or fetotoxic effect of vitamin D. As a result, vitamin D may be prescribed during pregnancy if needed. If necessary, vitamin D may be prescribed during breastfeeding.

(This card focuses on major safety information for medicinal products in order to minimize possible side effects that arise from improper use of medicinal products).

Pharmacovigilance Department

CONTACT DETAILS OF THE QPPV:

Phone: (962) (6) 5233349/113

email: raghad@sana-pharma.com

You can Also Report problems of Sana pharmaceutical products online via following link:

<http://www.sana-pharma.com/?q=node/3646>