

- Directions:

Ketofast[®] gel should be applied to the affected area 2-3 times a day for not more than 10 days.

Precautions:

- For external use only.
- Avoid the exposure of Ketofast[®] gel to mucous membrane areas (nose, ear, mouth, lip and genital area).
- Don't apply ketofast[®] gel to broken or infected skin.
- Avoid repetitive and excessive use of ketofast[®] gel.
- Ketofast[®] gel should be used with caution in patients with reduced liver, heart or kidney function.
- Avoid sun exposure for treated area during the treatment and 2 weeks after its discontinuation.
- Not recommended to use in child under 12 years.

- Contraindications:

- Hypersensitivity to the active substance or to any of the excipients of Ketofast[®] gel.
- History of any photosensitivity reaction or skin allergy to ketoprofen, tiaprofenic acid, fenofibrate or UV blocker or perfumes.
- Known hypersensitivity reactions, such as symptoms of asthma, allergic rhinitis or urticaria to fenofibrate, tiaprofenic acid, acetylsalicylic acid, or to other NSAIDs.

- Pregnancy and lactation:

- Ketofast[®] gel is contraindicated during the third trimester of pregnancy and should be avoided in first and second trimester.
- It isn't recommended to use Ketofast[®] gel in nursing mother.

(This card focuses on major safety information for medicinal products in order to minimize possible side effects that arise from improper use of medicinal products).

Pharmacovigilance Department

CONTACT DETAILS OF THE QPPV:

Phone: (962)/ (6) 5233349/113

email: e.alkhatatbeh@sana-pharma.com.

You can Also Report problems of Sana pharmaceutical products online via following link:

<http://www.sana-pharma.com/?q=node/3646>